



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ern Georgia, are intergrades, but those that I have seen are rather nearer the northern extreme than the Florida extreme, and I have therefore given the new name to the form inhabiting South Florida. Another reason that has induced me to do this is that the northern form never gets so far south as Miami in its winter migration while it is extremely common in winter in South Carolina; therefore the northern extreme can easily be got at the type locality of the species, while the Florida form stands apart by itself. The Museum of Comparative Zoölogy is fortunate in possessing a fine series, including both breeding and winter specimens of the Florida Bluebird, collected in 1871 by Messrs. Maynard and Henshaw, mostly at Miami — a point so far south that it represents the form in its extreme.

GENERAL NOTES.

Brünnich's Murre (*Uria lomvia*) at **Ottawa, Canada.** — On the 12th December, 1897, large numbers of this bird passed the city on the way South. The flight continued nearly the whole day. Quite a number of the birds were shot. — G. R. WHITE, *Ottawa, Ontario, Canada.*

Ross's Gull (*Rhodostethia rosea*) on **Bering Island.** — In my 'Ornith. Expl. Comm. Isls. and Kamtsch.' (1885), p. 315, I enumerated Ross's Gull among the birds of Kamchatka with some hesitation and without giving it a number since Saunders had queried the statement of Verreaux that the two specimens in the museum at Mayence actually came from that country. At the same time I pointed out that there was no improbability *per se* in the alleged locality being correct. I am now in position to affirm that this species occasionally straggles as far south on the Asiatic coast as Bering Island off the coast of Kamchatka. Last summer Mr. N. Grebnitski kindly presented me with a fully adult female of Ross's Gull obtained on Bering Island December 10, 1895. It is now in the U. S. National Museum, No. 162785. This is a very interesting addition to the avifauna of the Commander Islands. — LEONHARD STEJNEGER, *U. S. National Museum, Washington, D. C.*

The Scarlet Ibis — A Correction. — In 'The Auk,' XIV, 1897, p. 316, is a record by the present writer of the Scarlet Ibis taken in 1897 in the Arkansas Valley in Colorado. This was given on the authority of the